

Statistics Handbook

Contents	Page
Wilcoxon rank-sum test (Mann-Whitney equivalent)	2
Wilcoxon matched-pairs test	3
Normal Distribution	4
Z-test	
Related samples t-test	5
Unrelated samples t-test	6
Variance test (F-ratio)	7
Binomial distribution	8
Chi-square test	8
N-by-1 table	
Contingency table	
Correlation	10
Pearson's product-moment correlation	
Spearman's rank-order correlation	
Linear Regression	11
Box plot	12
Tables	Page
Critical values of Wilcoxon W (rank-sum)	2
Critical values of Wilcoxon T (matched-pairs)	3
Z distribution	4
Critical values of t	5
Critical values of F	7
Critical values of <i>Chi-square</i>	9
Critical values of r and r_s	11

Copyright (c) 2006. Andrew Wills. All rights reserved. **Version: 1.1.**

Reproduction is authorised without consent or fee only under the following terms: 1) All reproductions must include the above copyright statement and these terms. 2) Unlimited copies for non-profit educational institutions. 3) Unlimited copies for personal, non-commercial use.

All statistical tables were computed by the author.

Wilcoxon Rank-Sum Test

(Equivalent to the Mann-Whitney U test, except quicker, and the table has different values)

1. Rank all data, irrespective of group.
2. Calculate the sum of the ranks for the group with lower n . (If groups are of equal n , calculate the sum of the ranks for each group, and take the smaller).
3. The result is significant if your number is smaller or equal to the appropriate value in the tables below ($n_1 = n$ for smaller group, $n_2 = n$ for larger group).

Significance level: 0.05 one-tailed, 0.1 two-tailed

		n2														
		6	7	8	9	10	11	12	13	14	15	16	17	18	19	20
n1	6	28	30	32	34	35	37	39	41	43	44	46	48	50	52	53
	7		39	41	43	46	48	50	52	54	57	59	61	63	66	68
	8			52	54	57	60	62	65	67	70	73	75	78	81	83
	9				66	69	72	75	78	81	84	87	90	94	97	100
	10					83	86	90	93	96	100	103	107	110	114	117
	11						101	105	109	112	116	120	124	128	132	136
	12							121	125	130	134	138	142	147	151	155
	13								143	148	152	157	162	166	171	176
	14									167	172	177	182	187	192	197
	15										192	198	203	209	215	220
	16											220	226	232	238	244
	17												249	256	262	269
	18													281	287	294
	19														314	321
	20															349

Significance level: 0.025 one-tailed. 0.05 two-tailed

		n2														
		6	7	8	9	10	11	12	13	14	15	16	17	18	19	20
n1	6	26	28	29	31	32	34	36	37	39	40	42	44	45	47	48
	7		37	39	40	42	44	46	48	50	52	54	56	58	60	62
	8			49	51	53	56	58	60	63	65	67	70	72	75	77
	9				63	65	68	71	74	76	79	82	85	87	90	93
	10					79	82	85	88	91	94	97	100	104	107	110
	11						96	100	103	107	110	114	117	121	124	128
	12							116	119	123	127	131	135	139	143	147
	13								137	141	145	150	154	159	163	167
	14									160	165	169	174	179	184	188
	15										185	190	195	200	205	211
	16											211	217	223	228	234
	17												240	246	252	258
	18													271	277	283
	19														303	310
	20															337

Wilcoxon Matched-Pairs Test

1. Calculate the difference between each pair.
2. Remove pairs whose difference is zero, and reduce n accordingly.
3. Rank the differences of remaining pairs, ignoring their sign.
4. Calculate the sum of the ranks of the positive differences (T^+).
5. Calculate the sum of the ranks of the negative differences (T^-).
6. Let T be the smaller of T^+ and T^- .
7. The result is significant if T is smaller or equal to the appropriate value in the table below.

N	Significance	
	0.05 one-tailed (0.10 two-tailed)	0.025 one-tailed (0.05 two-tailed)
6	2	1
7	4	2
8	6	4
9	8	5
10	11	8
11	14	10
12	18	14
13	21	17
14	26	21
15	31	25
16	36	30
17	41	35
18	47	40
19	54	46
20	60	52

Normal distribution

Z-test

To find the probability with which a score X comes from a normal distribution with a mean of μ and a standard deviation of σ .

1. Calculate:

$$z = \frac{X - \mu}{\sigma}$$

2. Ignore the sign of z . The table below gives the one-tailed probability. Double the p values in the table for two-tailed probability.

Z-table

z	p	z	p	z	p	z	p
0.00	0.5000	1.52	0.0643	2.16	0.0154	2.76	0.0029
0.05	0.4801	1.54	0.0618	2.18	0.0146	2.78	0.0027
0.10	0.4602	1.56	0.0594	2.20	0.0139	2.80	0.0026
0.15	0.4404	1.58	0.0571	2.22	0.0132	2.82	0.0024
0.20	0.4207	1.60	0.0548	2.24	0.0125	2.84	0.0023
0.25	0.4013	1.62	0.0526	2.26	0.0119	2.86	0.0021
0.30	0.3821	1.64	0.0505	2.28	0.0113	2.88	0.0020
0.35	0.3632	1.66	0.0485	2.30	0.0107	2.90	0.0019
0.40	0.3446	1.68	0.0465	2.32	0.0102	2.92	0.0018
0.45	0.3264	1.70	0.0446	2.34	0.0096	2.94	0.0016
0.50	0.3085	1.72	0.0427	2.36	0.0091	2.96	0.0015
0.55	0.2912	1.74	0.0409	2.38	0.0087	2.98	0.0014
0.60	0.2743	1.76	0.0392	2.40	0.0082	3.00	0.0013
0.65	0.2578	1.78	0.0375	2.42	0.0078	3.02	0.0013
0.70	0.2420	1.80	0.0359	2.44	0.0073	3.04	0.0012
0.75	0.2266	1.82	0.0344	2.46	0.0069	3.06	0.0011
0.80	0.2119	1.84	0.0329	2.48	0.0066	3.08	0.0010
0.85	0.1977	1.86	0.0314	2.50	0.0062	3.10	0.0010
0.90	0.1841	1.90	0.0287	2.52	0.0059	3.12	0.0009
1.00	0.1587	1.92	0.0274	2.54	0.0055	3.14	0.0008
1.05	0.1469	1.94	0.0262	2.56	0.0052	3.16	0.0008
1.10	0.1357	1.96	0.0250	2.58	0.0049	3.18	0.0007
1.15	0.1251	2.00	0.0228	2.60	0.0047	3.20	0.0007
1.20	0.1151	2.02	0.0217	2.62	0.0044	3.22	0.0006
1.25	0.1056	2.04	0.0207	2.64	0.0041	3.24	0.0006
1.30	0.0968	2.06	0.0197	2.66	0.0039	3.26	0.0006
1.35	0.0885	2.08	0.0188	2.68	0.0037	3.30	0.0005
1.40	0.0808	2.10	0.0179	2.70	0.0035	3.50	0.0002
1.45	0.0735	2.12	0.0170	2.72	0.0033	3.75	0.0001
1.50	0.0668	2.14	0.0162	2.74	0.0031	4.00	0.0000

Related samples t-test

1. Calculate the difference (D) between each pair.
2. Calculate the mean of the differences, \bar{D} .
3. Calculate the standard deviation of the differences

$$s_D = \sqrt{\frac{\sum (D - \bar{D})^2}{N - 1}}$$

4. Calculate the standard error of the differences:

$$s_{\bar{D}} = \frac{s_D}{\sqrt{N}}$$

5. Calculate the t statistic:

$$t = \frac{\bar{D}}{s_{\bar{D}}}$$

6. Ignore the sign of t . The result is significant if t is greater than the appropriate value in the t -table.

- For a related samples t -test with N pairs, $df = N - 1$.

t-table

df	Significance Level			
	0.1 two-tailed (0.05 one-tailed)	0.05 two-tailed (0.025 one-tailed)	0.025 two-tailed (0.0125 one-tailed)	0.01 two-tailed (0.005 one-tailed)
1	6.314	12.706	25.452	63.656
2	2.920	4.303	6.205	9.925
3	2.353	3.182	4.177	5.841
4	2.132	2.776	3.495	4.604
5	2.015	2.571	3.163	4.032
6	1.943	2.447	2.969	3.707
7	1.895	2.365	2.841	3.499
8	1.860	2.306	2.752	3.355
9	1.833	2.262	2.685	3.250
10	1.812	2.228	2.634	3.169
11	1.796	2.201	2.593	3.106
12	1.782	2.179	2.560	3.055
13	1.771	2.160	2.533	3.012
14	1.761	2.145	2.510	2.977
15	1.753	2.131	2.490	2.947
16	1.746	2.120	2.473	2.921
17	1.740	2.110	2.458	2.898
18	1.734	2.101	2.445	2.878
19	1.729	2.093	2.433	2.861
20	1.725	2.086	2.423	2.845

Unrelated samples t-test

Equal N

1. Let N be the sample size of each group.
2. Calculate the mean for each group, \bar{X}_1 and \bar{X}_2
3. Calculate the variance for each group, s^2_1 and s^2_2 .

$$s^2 = \frac{\sum (X - \bar{X})^2}{N - 1}$$

4. Calculate the t statistic:

$$t = \frac{(\bar{X}_1 - \bar{X}_2)}{\sqrt{\frac{s^2_1 + s^2_2}{N}}}$$

5. Ignore the sign of t . The result is significant if t is greater than the appropriate value in the t -table.
 - $df = 2N - 2$.

Unequal N

1. Let N_1 and N_2 be the sample sizes of the two groups.
2. Calculate the mean for each group, \bar{X}_1 and \bar{X}_2
3. Calculate the variance for each group, s^2_1 and s^2_2 .

$$s^2 = \frac{\sum (X - \bar{X})^2}{N - 1}$$

4. Calculate the pooled variance estimate:

$$s^2_p = \frac{(N_1 - 1)s^2_1 + (N_2 - 1)s^2_2}{N_1 + N_2 - 2}$$

5. Calculate the t -statistic:

$$t = \frac{\bar{X}_1 - \bar{X}_2}{\sqrt{s^2_p \left(\frac{1}{N_1} + \frac{1}{N_2} \right)}}$$

6. Ignore the sign of t . The result is significant if t is greater than the appropriate value in the t -table.
 - $df = N_1 + N_2 - 2$

Variance test

1. Calculate the variance for each of the two groups:

$$s^2 = \frac{\sum(X - \bar{X})^2}{N - 1}$$

2. Let S_L^2 be the smaller of the two variances, and S_H^2 be the larger.
3. Calculate the F-ratio:

$$F = \frac{S_H^2}{S_L^2}$$

4. The variances are significantly different if F is greater than the appropriate value in the F table.
 - The degrees of freedom for the numerator are $(N_H - 1)$, where N_H is the sample size for the group with higher variance. df for the denominator are $(N_L - 1)$.
 - This is a two-tailed test.

F table

Degrees of Freedom for Denominator	Significance Level: 0.025 one-tailed, 0.05 two-tailed											
	Degrees of Freedom for Numerator											
	1	2	3	4	5	6	7	8	9	10	15	20
1	647.8	799.5	864.2	899.6	921.8	937.1	948.2	956.6	963.3	968.6	984.9	993.1
2	38.51	39.00	39.17	39.25	39.30	39.33	39.36	39.37	39.39	39.40	39.43	39.45
3	17.44	16.04	15.44	15.10	14.88	14.73	14.62	14.54	14.47	14.42	14.25	14.17
4	12.22	10.65	9.98	9.60	9.36	9.20	9.07	8.98	8.90	8.84	8.66	8.56
5	10.01	8.43	7.76	7.39	7.15	6.98	6.85	6.76	6.68	6.62	6.43	6.33
6	8.81	7.26	6.60	6.23	5.99	5.82	5.70	5.60	5.52	5.46	5.27	5.17
7	8.07	6.54	5.89	5.52	5.29	5.12	4.99	4.90	4.82	4.76	4.57	4.47
8	7.57	6.06	5.42	5.05	4.82	4.65	4.53	4.43	4.36	4.30	4.10	4.00
9	7.21	5.71	5.08	4.72	4.48	4.32	4.20	4.10	4.03	3.96	3.77	3.67
10	6.94	5.46	4.83	4.47	4.24	4.07	3.95	3.85	3.78	3.72	3.52	3.42
11	6.72	5.26	4.63	4.28	4.04	3.88	3.76	3.66	3.59	3.53	3.33	3.23
12	6.55	5.10	4.47	4.12	3.89	3.73	3.61	3.51	3.44	3.37	3.18	3.07
13	6.41	4.97	4.35	4.00	3.77	3.60	3.48	3.39	3.31	3.25	3.05	2.95
14	6.30	4.86	4.24	3.89	3.66	3.50	3.38	3.29	3.21	3.15	2.95	2.84
15	6.20	4.77	4.15	3.80	3.58	3.41	3.29	3.20	3.12	3.06	2.86	2.76
16	6.12	4.69	4.08	3.73	3.50	3.34	3.22	3.12	3.05	2.99	2.79	2.68
17	6.04	4.62	4.01	3.66	3.44	3.28	3.16	3.06	2.98	2.92	2.72	2.62
18	5.98	4.56	3.95	3.61	3.38	3.22	3.10	3.01	2.93	2.87	2.67	2.56
19	5.92	4.51	3.90	3.56	3.33	3.17	3.05	2.96	2.88	2.82	2.62	2.51
20	5.87	4.46	3.86	3.51	3.29	3.13	3.01	2.91	2.84	2.77	2.57	2.46

Binomial test

Let the outcomes be P and Q , and the number of trials be N . The probability of the P outcome occurring *exactly* X times is given by:

$$\text{prob}(X) = \frac{N!}{X!(N-X)!} p^X q^{(N-X)}$$

where

$$\begin{aligned} p &= \text{The probability of the } P \text{ outcome} \\ q &= \text{The probability of the } Q \text{ outcome} \\ p+q &= 1 \end{aligned}$$

$N!$ is **N factorial** e.g. $4! = 4 \times 3 \times 2 \times 1 = 24$.

Chi-square test

- The formula for chi-square is:

$$\chi^2 = \sum \frac{(O-E)^2}{E}$$

where O = observed frequency and E = expected frequency. Expected values depend what you're trying to do. Some examples:

N-by-1 table

There is one variable, with N levels. There are N observed frequencies, and the null hypothesis is that they do not differ. Expected frequency is the mean of the observed frequencies. Degrees of freedom = $N - 1$.

Example:

	Heads	Tails
Observed	83	17
Expected	50	50

Contingency table

There are two variables. One variable has M levels. The other variable has N levels. There are $N \times M$ observed frequencies. The null hypothesis is that the two variables are independent.

Expected frequency = Row Total x Column Total / Grand Total

Degrees of freedom = $(N-1)(M-1)$

Example (expected frequencies given in brackets)

	Non-smoker	Smoker	TOTAL	
Male	36 (26)	42 (52)	78	
Female	23 (33)	76 (66)	99	
TOTAL	59	118	177	←Grand total

Test of significance

- The result is significant if the calculated value of chi-square exceeds the appropriate value on the chi-square table.

Chi-square table

df	Significance Level			
	0.1	0.05	0.025	0.01
1	2.706	3.841	5.024	6.635
2	4.605	5.991	7.378	9.210
3	6.251	7.815	9.348	11.345
4	7.779	9.488	11.143	13.277
5	9.236	11.070	12.832	15.086
6	10.645	12.592	14.449	16.812
7	12.017	14.067	16.013	18.475
8	13.362	15.507	17.535	20.090
9	14.684	16.919	19.023	21.666
10	15.987	18.307	20.483	23.209
11	17.275	19.675	21.920	24.725
12	18.549	21.026	23.337	26.217
13	19.812	22.362	24.736	27.688
14	21.064	23.685	26.119	29.141
15	22.307	24.996	27.488	30.578
16	23.542	26.296	28.845	32.000
17	24.769	27.587	30.191	33.409
18	25.989	28.869	31.526	34.805
19	27.204	30.144	32.852	36.191
20	28.412	31.410	34.170	37.566

Correlation

Calculation of the correlation co-efficient r between two variables x and y
(*Pearson product-moment correlation*)

1. Calculate the mean of each variable, \bar{X} and \bar{Y} .
2. For each point, calculate $(X - \bar{X})$, $(X - \bar{X})^2$, $(Y - \bar{Y})$ and $(Y - \bar{Y})^2$
3. From these values, calculate the standard deviation of each variable

$$s = \sqrt{\frac{\sum (X - \bar{X})^2}{N - 1}}$$

4. Also from the values calculated in 2), calculate the *co-variance* of x and y :

$$\text{COV}_{XY} = \frac{\sum (X - \bar{X})(Y - \bar{Y})}{N - 1}$$

5. Calculate r

$$r = \frac{\text{COV}_{XY}}{s_X s_Y}$$

6. The correlation is significant if r exceeds the appropriate value on the table below. As a check of your arithmetic, note that r always lies between -1 and +1.
 - Spearman's rank-order correlation co-efficient r_s can be calculated by applying the above procedure to the ranks of x and y , instead of the raw scores. **Rank x and y separately.**
 - In the context of correlation, "tails" refers to the sign of r . If the test is for r simply being different from zero, then the test is two-tailed.

Linear Regression

Procedure for finding the best-fitting line of the form

$$y = \mathbf{b}.x + \mathbf{a}$$

1. Calculate the gradient of the best-fitting straight line from the co-variance of x and y , and the variance of x .

$$b = \frac{\text{COV}_{XY}}{s_x^2}$$

2. Calculate the intercept of that line:

$$a = \bar{Y} - b\bar{X}$$


3. If the correlation co-efficient r is significantly different from zero, then b is significantly different from zero.

Critical Values of r (and r_s)

N	Two-tailed significance Level		
	<i>0.1 two-tailed</i> 0.05 one-tailed	<i>0.05 two-tailed</i> 0.025 one-tailed	<i>0.01 two-tailed</i> 0.005 one-tailed
9	0.575	0.657	0.785
10	0.544	0.625	0.754
11	0.517	0.596	0.726
12	0.494	0.571	0.701
13	0.473	0.549	0.677
14	0.455	0.529	0.656
15	0.439	0.511	0.637
16	0.424	0.495	0.619
17	0.411	0.480	0.602
18	0.399	0.467	0.587
19	0.388	0.454	0.572
20	0.377	0.442	0.559

The values given in the table are based on an approximation that is accurate to within 0.02 over the range covered.

Box Plot



Place your data in rank order, lowest to the left. The equations below give the *position* of the median etc. counting from the left. The median is given by the *number* at that position. If the position is fractional (e.g. 5.5), average the numbers at the two whole-number positions (e.g. at positions 5 and 6).

Median position = $(N+1) / 2$

Lower hinge position* = $(\text{median position} + 1) / 2$

Upper hinge position = $N + 1 - \text{lower hinge position}$

Inter-quartile range (IQR) : Difference between data values at upper and lower hinge positions

Whisker = $1.5 \times \text{IQR}$

Outliers: Typically, points more than two whiskers from the nearest hinge.

* *Ignore fractional component of median position in calculation of lower hinge position.*